



VELDGAINER 95

PROTEIN MINERAL TRACE MINERAL VITAMIN CONCENTRATE FOR HOME MIXING OF VELD SUPPLEMENTS FOR GROWING OUT AND FINISHING OF LIVESTOCK ON VELD PASTURE

KEY BENEFITS

- Highly concentrated – make maximal use of own / local feed ingredients
- Supplies a moderate level of NPN - suitable for all ruminants on natural veld throughout the year
- Contains NPN in correct ratio with sulphur for optimal microbial utilization of grazing
- Contains high levels of phosphate and magnesium for maximal growth
- Contains an optimal level of monensin – stimulates intake and digestion of veld and improves growth

EXAMPLES OF SUPPLEMENTS

INGREDIENTS	VELD STARTER LICK		VELD GROWER LICK		VELD FINISHER LICK	
Grain meal / Hominy Chop	700	675	775	750	850	825
Soy oilcake meal 47	50		50		50	
Cotton oilcake meal 38		75		75		75
Salt	150	150	100	100	50	50
Veldgainer 95	100	100	75	75	50	50
Total (kg)	1000	1000	1000	1000	1000	1000
Composition (%)						
Protein	17.1	17.4	15.3	15.6	13.5	13.8
Protein ex NPN (% of total)	55.5	54.6	46.6	45.7	35.2	34.4
Energy (MJ ME/kg)	9.3	9.2	10.2	10.1	11.2	11.1
Calcium	1.67	1.67	1.26	1.26	0.85	0.85
Phosphorus	0.73	0.76	0.62	0.65	0.51	0.55
Sulphur	0.41	0.42	0.35	0.35	0.28	0.28
Magnesium	0.34	0.36	0.30	0.32	0.25	0.27
Intake – Large stock (g/day)	1000 - 2000		1500 - 3000		2000 - 4000	
Intake – Small stock (g/day)	200- 400		300 - 600		400 - 800	

FEEDING RECOMMENDATIONS

- Replace up to 10% of the grain meal with extra salt for the first 7 days on the starter lick for adaptation
- Replace soybean- or cotton oilcake meal with 100 kg of sunflower- or canola oilcake meal per mixture if preferred
- After adaptation the best regulated intakes will be achieved by providing these licks unrestricted at all times
- Provide adequate trough space at all times to ensure adequate intake by all animals
- Consult a Feedtek nutritionist if intakes differ widely from recommended intakes after a period of 14 days
- Provide grazing and / or roughage unrestricted together with these licks at all times
- Replace up to 10% of the grain meal with molasses meal or molasses syrup if preferred
- These licks contain NPN and a medicament – mix thoroughly, protect against rain and do not feed to horses
- Provide the starter lick for at least 14 days before moving to the grower lick
- Move over to the finisher lick as soon as animals have grown out in frame for the final finishing phase
- For preparing animals for shows and auctions it is recommended that they remain on the grower lick until the end of the feeding period

